17.1. Jod weeg

5.

The British Society for Parasitology -Spring Meeting, Canterbury, 5th - 7th April 1978.

APPLICATION TO DISPLAY A FOSTER (maximum dimensions 2 metres wide by 1.5 metres high)

Applications received after 23rd January will not be accepted. Please read enclosed notes on the seeting.

- 1. AUTHOR (S) Drs. M.H. MIRCK (Give title: Prof.Dr. etc)
- 2. PLACE OF RESEARCH Department of Veterinary Medicine.

Veterinary Faculty, State University of Utrecht. 'ne Welherland's

- 3. TELEPHONE NUMBER __030-531094
- 4. TITLE OF POSTER

 (Please be as informative and succinct as possible. Give generic and specific names in full)

Parasitological faecal examination of 3791 paarden on

ponies in the Netherlands.

5. CONTENT OF POSTER

(Brief but clear statement on content of poster, not more than 100 words. This information is required to help the organisers plan the poster displays).

Parasitological faecal examination was carried out of 3791 horses and ponies brought to the Department of Veterinary Medicine of the State University of Utrecht from 1973 up to 1977 inclusive, because of different motives.

The results were arranged according to the age of the patient and also of the month of arrival. Generally it appears that 62,3 percent of the faecal samples obtained worm-eggs. In 57,3 percent there were strongyle-eggs present, in 6,1 percent eggs of Parascaris equorum, in 1,2 percent eggs of Oxyuris equi, in 0,2 percent eggs/larvae of Dictyocaulus arrifieldi, in 0,6 percent eggs of Fasciola hepatica, in 4,7 percent eggs of Strongy-loides westeri (only in horses and ponies under one year of age), in 2,5 percent eggs of Anoplocephala spp. and in 0,3 percent *gocysts of PLEASE REPEAT THE TITLE OF YOUR POSTER IN THE SPACE ALLOCATED BELOW

6. PLEASE REPEAT THE TITLE OF YOUR POSTER IN THE SPACE ALLOCATED BELO AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE FOX PROVIDED.

Parasitological faecal examination of 3791 horses and ponies in the Netherlands.

NAME

ADDRESS

drs. M.H. Mirck,
Department of Veterinary Medicine,
State University of Utrecht,
Yalelaan 16,
"De Uithof"
Utrecht,
The Netherlands.

APPLICATION TO DISPLAY A POSTER (maximum dimensions 2 metres wide by 1.5 metres high)

Applications received after 23rd January will not be accepted. Please read enclosed notes on the meeting.

- 1. AUPHOR (S) Dr.L. Gustowska, Dr. E.J. Ruitenberg and Anneke Elgersma (Give title: Prof. Dr. etc)
- 2. PLACE OF RESEARCH National Institute of Public Health, Bilthoven, The Nether-Laboratory for Pathology
- 3. TELEPHONE NUMBER 030 789111 ext. 1075
- 4. TITLE OF POSTER
 (Please be as informative and succinct as possible. Give generic and specific names in full)

Immunopathological studies of the muscle phase of Trichinella

spiralis; a comparison between normal and congenitally athymic (nude)

5. CONTENT OF POSTER

(Brief but clear statement on content of poster, not more than 100 words. This information is required to help the organisers plan the poster displays).

Study of the immunopathology (cellular infiltrate) of both intestinal and muscle phase of <u>Trichinella spiralis</u> in congenitally athymic (nude) nu/nu mice compared to their heterozygous (thymus-containing) (+/nu) littermates. Basophilic transformation of muscle cells (i.e. differentiation to nurse cell) was T-independent, whereas cellular infiltrate around the encysted larva was mainly T-dependent.

Basophilic transformation preceded the presence of muscle larvae in both $+/\underline{n}\underline{u}$ and $\underline{n}\underline{u}/\underline{n}\underline{u}$ mice. Muscle larvae were observed earlier in the tongue of $+/\underline{n}\underline{u}$ mice. Later in the infection the number of muscle larvae in the $\underline{n}\underline{u}/\underline{n}\underline{u}$ mice exceeded that of the $+/\underline{n}\underline{u}$ mice.

Antibody production (measured with IF, ELISA, PCA) was only observed in infected +/nu mice. The interdependence of the results is discussed.

6. HLEASE REPEAT THE TITLE OF YOUR POSTER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE POX PROVIDED.

NAME

ADDRESS

Dr. E.J.Ruitenberg

National Institute of Public Health P.O. Box 1 Bilthoven The Netherlands

TITIE: Immunopathological studies of the muscle phase of <u>Trichinella</u>

<u>spiralis</u>; a comparison between normal and congenitally athymic

the first of F. Lity for Paralitology -Spring Meeting, Canterbury,

1.5 netres high)

I to end notes on the secting. And danuary will not be accepted. Please

. All hok . Drs.J.F.Sluiters (Give tille. Prof. Dr. etc.)

.. H.ACE OF RESEARCH Free University, Laboratory of Medical Parasitology,

v.d. Boechorststraat 7, Amsterdam, The Netherlands.

5.

TELEPHONE NUMBER 020 - 5483979

TITLE OF ROSPER

(Mease be as informative and succinct as possible. Give generic

The development of the gonad of Lymnaea stagnalis during an infection

with Trichobilharzia ocellata.

CONTENT OF POSTER

(Brief but clear statement on content of poster, not more than 100 word... This information is required to help the organisers plan

Juvenile L. stagnalis are heavily infected with T. ocellata. In the presence of the primary sporocysts already the growth of the ovotestis is strongly retarded. The total number of gametes remains small compared to the controls. Preliminary results indicate that during development of the secondary sporocysts and after patency is reached sperms are found in the spermoviduct and oocytes mature, but in low numbers. Ovulation, however, seems not to occur: oocytes degenerate and oviposition is not observed. These data suggest an influence of the parasite on the endocrine control of reproduction of the host.

PLEASE REPEAT THE TITLE OF YOUR POSTER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE HOX PROVIDED.

NAME

ADDRESS

Drs.J.F.Sluiters

Free University, Laboratory of Medical Parasitology, v.d.Boechorststraat 7, Amsterdam, The Netherlands.

TITLE: The development of the gonad of Lymnaea stagnalis during an infection with Trichobilharzia ocellata.

31.77

the british So rety for Parasitology Spring Meeting, Canterbury, 111. 7th April 1975.

(maximum dimensions 2 metres wide by 1.5 metres high)

read. lored notes on the seeting.

- . AU HOR (E) Drs. J.J.Mellink (Give title: Prof.Dr. (tc)
- Free University Amsterdam THE NETHERLANDS
- TELEPHONE NUMBER 020-5483981
- (Mease be as informative and succinct as possible. Give generic and specific names in full)

Vector - host relationships in haematophagy by the yellow fever mosquito, Aedes aegypti (L.).

(Brief but clear statement on content of poster, not more than 100 the poster displays)

In our laboratory mosquito blood-feeding is being studied in relation to disease transmission. Detailed observations were made of the feeding process and special attention was paid to possible sources of variations in mosquito feeding behaviour, also related to host reactions. Functional aspects of mosquito salivation were elucidated by comparing salivating and non-salivating mosquitoes. Research on the potential relevance of variations in mosquito feeding behaviour for viral transmission is in progress. Preliminary results of the latter will be included in the poster presentation.

AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE HOX PROVIDED.

Wester - host relationships in haematophagy by the yellow fever

mosquito, Aedes aegypti (L.)

NAME

Drs. J.J.Mellink

ADDRESS

Laboratory of Medical Parasitology Free University

P.O.Box 7161

Amsterdam.

THE NETHERLANDS

the present Society for Paramitology - Spring Meeting, Canterbury.

AL ANTHORY OF PRESENT A FIRE PAPER (15 mins. duration inclusive of discussion)

- t. a' History or Hole (,) Dr.F.R.Opperdoes (ve title: Prof, Dr. etc.)
- C. 1. NOE OF RESEARCH Section for Medical Enzymology and Molecular Biology, University of Amsterdam, Eerste Constantijn Huygensstraat 20, Amsterdam The Nather lands a
- 020 181313
- 4. TITLE OF PAPER

(please be as informative and succinct as possible. Give generic and specific names in full)

The glycosome of trypanosomes.

RESUME OF PAPER

(brief but clear resume of content of paper, not more than 100 words. This will be used to hel; the organisers but will not be published)

The trypomastigote of Trypanosoma brucei is completely dependent on glycolysis for its energy production. This glycolytic pathway is modified in several res-

- A. Glucose is converted into pyruvate rather than lactate.
- B. NADH produced in glycolysis is reoxidized indirectly by 0 via a glycerol-3-phosphate: dihydroxyacetone phosphate cycle.

C. Dismutation of glucose into pyruvate and glycerol under anaerobic conditions gives rise to ATP synthesis.

Fractionation of homogenates of T.brucei shows that 9 glycolytic enzymes involved in the conversion of glucose and glycerol into 3-P-glycerate are particle bound and show latency, whereas in isopycnic centrifugation these enzymes all band together at a density of 1.23 g/cm3. Electron microscopic examination shows that this part of the gradient is highly enriched in microbody-like organelles. These observations indicate that part of the glycolytic pathway in T.brucei is located inside a microbody. Because this microbody lacks any peroxisomal or microbody marker enzyme tested thusfar, we have proposed the name "glycosome" (For organiser's use)

SESSI ON			
MOTORICA	TIME	DACTOR	
	-	DATE	

PLEASE REPEAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

The glycosome of trypanosomes

VAME

ADDRESS

Dr.F.R.Opperdoes

Section for Medical Enzymology and Molecular Biology, University of Amsterdam, Eerste Constantijn Huygensstraat 20, The Netherlands.

for this organelle.

Preliminary experiments, with identical results, with the distantly related insect trypanosome Crithidia luciliae suggest that location of glycolysis inside a glycosome is a general property of trypanosomes.

The function of the glycosome and its involvement in aerobic and anaerobic glycolysis will be discussed.

The british Society for Parasitology	- Spring	Meeting, Canterbury.
AL ACTION TO PRESEND A DIEGO		

- : 4 1 received after corn January w 11 not be accepte. Please read enotes d notes on the meet ng.
- 1. AFPHOR (:) Drs.J.H.J.Hoeymakers, F.Fase-Fowler and Prof.Dr.P.Borst (ive title: Prof, Dr. etc.)
- In ACE OF RESEARCH University of Amsterdam, Dept.of Medical Enzymology, Jan Swammerdam Institute, 1e Const. Huygensstr. 20,
- TELEPHONE NUMBER Amsterdam, The Netherlands
- 020-181313
- TITLE OF PAPER (please be as informative and succinct as possible. Give generic Studies on KINETOPLAST DNA in trypanosomes.

5. RESUME OF PAPER

(brief but clear resume of content of paper, not more than 100 words. This will be used to hel; the organisers but will not be published)

We shall discuss our recent experiments on:

- genetic complexity of kinetoplast DNA (kDNA)
- sequence evolution of mini- and maxicircles in Crithidia
- structure of kDNA networks of Crithidia, Trypanosoma brucei, T.equiperdum, and T.evansi.
- structure of putative intermediates in kDNA replication.
- morphology of kDNA in gently lysed cells.

t. tFor organi	SESSION		
15. 45		TIME	DATE
		OUR PAPER IN THE SPACE	ALLOCATED BELOW BOX PROVIDED.
		NA in trypanosomes	
VAME	Drs.H.J.Hoe	eymakers eymakers	
ADDRESS	University	of Amsterdam,	

Dept.of Medical Enzymology, Jan Swannerdam Institute, 1e Const. Huygensstr. 20, Amsterdam, The Netherlands.

API_JCATION TO PRESENT A FIRE PAPER (15 mins. duration inclusive of discussion)

Paper received after Bra January w 11 not be accepte. Please read enclored notes on the meet ng.

- 1. At PhoR (:) Dr. W. Eling and Drs. A. van Zon (tive title: Prof, Dr. etc.)
 - Department of Cytology and Histology
 University of Nijmegen, The Netherlands
- 3. TELEPHONE NUMBER 080-514665
- 4. TITLE OF PAPER (please be as informative and succinct as possible. Give generic and specific names in full)

Corticosteroids and malaria (Plasmodium berghei) immunity in mice.

5. RESUME OF PAPER

(brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be published)

Pregnancy induced immunodepression was observed in experimental malaria. A high proportion of immune mice experienced a frequently fatal breakthrough of malaria during pregnancy. A correlation between changes in serum corticosteroid levels and breakthrough of malaria during pregnancy initiated studies on the effect of these hormones on the malaria immune response in non-pregnant mice. The effect of corticosteroids on an established immunity was slight and complex. The strongest effect was observed during sensitization. Corticosteroid sensitive cells are apparently not important during early antigen recognition but functions in a later phase of the sensitization period. The mechanics of the immune response appear to change, however, after the challenge infection.

t. a	(For_	organi	ser.	s	use)

SESSION	mr		
- DOULOIT	TIME	DACRE	
		 DATE	
		1910 J. S. C.	

PLEASE REPEAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

Corticosteroids and malaria (Plasmodium berghei) immunity in mice.

.VAME

Dr. W. Eling and Drs. A. v. Zon

ADDRESS

Department of Cytology and Histology University of Nijmegen The Netherlands 3.

3. The British S recty for Parasitology - Spring Meeting, Canterbury. 5th - 7th Apr 1 100. APPLICATION TO PRESENT A FIRE PAPER (15 mins, duration inclusive of discussion) Paper received after Sre January w 11 not be accepter. Please read encloand notes on the meet ag. 1. AT PHOR (S) Drs. A. van Zon and Dr. W. Eling (Live title: Prof, Dr. etc.)
Department of Cytology and Histology FLACE OF RESEARCH University of Nijmegen, The Netherlands TELEPHONE NUMBER 080-514664 4. TITLE OF PAPER (please be as informative and succinct as possible. Give generic and specific names in full) Pregnancy and immunity in the Plasmodium berghei - mouse model RESUME OF PAPER (brief but clear resume of contert of paper, not more than 100 words. This will be used to hel: the organisers but will not be published) In the P. berghei-mouse model a high proportion of previously immune mice experienced a frequently fatal breakthrough of malaria during pregnancy. Due to the state of premunition associated with immunity in this model a pregnancy induced immunodepression can easily be observed. Not all immune mice with persisting parasites experienced a breakthrough during pregnancy. Such pregnancy associated recrudescences were not found to be associated particularly with the first pregnancy in contrast to what has been reported for human malaria. Chemoprophylaxis during the first pregnancy followed by reinfection did not protect against a relapse of malaria during a subsequent pregnancy. (, (For organiser's use) SESSION TIME PLEASE REPEAR THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN IN. BOX PROVIDED

FTLF Pregnancy and immunity in the Plasmodium berghei - mouse model

.VAME

Drs. A. van Zon and Dr. W. Eling

ADDRESS

Department of Cytology and Histology University of Nijmegen The Netherlands

APPLICATION TO PRESENT A FREE PAPER (15 mins. duration inclusive of discussion)

Papers received after 23rd January will not be accepted. Please read enclosed notes on the meeting.

- 1. AUTHOR (S) Dr. J.P. Verhave, Drs. A. Vermeulen, Dr. J.H.E.T. Meuwissen (Give title: Prof, Dr. etc.)
- 2. PLACE OF RESEARCH Inst. Med. Parasitology, University of Nijmegen, The Netherlands
- 3. TELEPHONE NUMBER 080-514306
- 4. TITLE OF PAPER

 (please be as informative and succinct as possible. Give generic and specific names in full)

 Specificity of anti-sporozoite antibody responses induced by Plasmodium berghei
- 5. RESUME OF PAPER
 (brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be published)

Chloroquinized rats, repeatedly exposed to sporozoites, show a high IFA-antibody response in their serum, when tested with sporozoite antigen (SPOR), but lower titres when tested with bloodform antigen (SCH) or exoerythrocytic forms (EEF). Adsorbtion with sporozoites leaves the anti-EEF, as well as the anti-SCH titres intact. Adsorbtion of such sera with freeze-thawed bloodform sediment reduces the anti-SCH titres and the anti-EEF titres, but it does not touch the anti-SPOR titres. Immunization of rats under protection of pyrimethamine or proguanil, of rabbits, or of rats with attenuated sporozoites excludes EEF-development; yet it induces low titres against EEF and SCH.

Apart from specific sporozoite determinants, there are common antigenic determinants present on all stages of the parasite.

6.	(For_	organi	ser'	s	use)	

SESSION TIME DATE

7. PLEASE REPEAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

TITLE Specificity of anti-sporozoite antibody responses induced by P. berghei

NAME

Dr. J.P. Verhave

ADDRESS

Inst.Med.Parasitology, Catholic University

Geert Grooteplein Z 24

NIJMEGEN

The British Senet, for Parasitology - Spring Meeting, Canterbury.

AT ACCUSTOM TO PRESENT A FILE PAPER (15 mins. duration inclusive of discussion)

: 4 1 not be accepte: Please read enclased notes on the meet ug.

- 1. At IHOK (DCH. J. van der Kaay, A.C.L. Sikkema and L. Boorsma (ive title: Prof, Dr. etc.)
- 2. H. MCE OF RESEARCH Leiden University, Medical Center, Department of Parasitology, Rapenburg 33, Leiden, the Netherlands TELEPHONE NUMBER
- 071-148333 ext 7442
- TITLE OF PAPER (please be as informative and succinct as possible. Give generic

Observations on the development of Plasmodium berghei berghei (ANKA) in a susceptible and refractive strain of Anopheles atroparvus van Thiel.

RESUME OF PAPER (brief but clear resume of content of paper, not more than 100 words. This will be used to hel; the organisers but will not be

The sporogony of P.berghei berghei is studied in a fully susceptible and a highly refractive strain of A.atroparvus. In the susceptible strain the production of ookinetes results in the development of a high number of oocysts. Though in the refractive strain ookinetes are produced in equally large numbers, they are apparently not able to pass a thin layer secreted by the midgut. Ookinetes are probably immobilized before this layer and seems to be barred from penetrating the epithelial lining of the mid-gut wall. Several of the observations are being discussed.

(For organiser's use)

SI	ESSION	TIME		
			-	DATE
HE	TITLE OF VOU			

PLEASE REPEAR THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

Observations on the development of Plasmodium berghei berghei (ANKA) in a susceptible and refractive strain of Anopheles atroparvus van Thiel.

VAME

Dr. H.J. van der Kaay

ADDRESS

Leiden University, Medical Center, Department of Parasitology, Rapenburg 33, Leiden, The Netherlands.

The British Swrety for Parabitology - Spring Meeting, Canterbury.

THOR TO PRESENT A FIEE PAPER (15 mins. duration inclusive of discussion)

rue) a note: on the meet ng.

- 1. A THUK () Dr. Wedad Tadros & Prof. Dr. J.J. Laarman
- 1. WE OF RESEARCH Dept. of Parasitology, University of Amsterdam. Institute of Tropical Hygiene, dpt. of the
- Royal Tropical Institute. The Netherlands TELEPHONE NUMBER 020 944422
- TITLE OF PAPER

(please be as informative and succinct as possible. Give generic and specific names in full)

CURRENT CONCEFTS ON THE TAXONOMY AND NOMENCLATURE OF TISSUE STAGE-FORMING DISPORIC TETRAZOIC EIMERIID COCCIDIA.

5. RESUME OF PAPER

(brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be

Controversial modern proposals and concepts on the taxonomy and nomenclature of this group: of coccidia will be briefly outlined. The pitfalls of using the morphology of the tissue stage in the intermediate host or the pattern of life cycle (homo- or heteroxenous) as main criterion for taxonomy will be discussed. The suthers present main criterion for taxonomy will be discussed. The authors present their own proposal for (i) emending the definition of the family Eimeriidae to encompass eimeriid coccidia with tissue stages in the same or intermediate hosts, (ii) re-erecting and re-defining the subfamily Eimeriinae for eimeriid coccidia shed unsporulated, the genera Toxoplasma, Hammondia and Besnoitia being synonymized with the genus Isospora; (iii) recognizing a separate subfamily for obligatorily heteroxenous eimeriid coccidia shed sporulated and lacking pre-

f	(For organiser's	use)

in white is the settled in a technical in most on the

SESSION	OTTAG		
	TIME	DATE	
HE PT IN DE LATE THE			

PLEASE REPEAU THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

ITHE Current concepts on the taxonomy and nomenclature of tissue stage forming disporic tetrazoic eimerlia coccidia.

VAME

ADDRESS

Dr. Wedad Tadros & Prof. Dr. J.J. Laarman, Institute of Tropical Hygiene, Mauritskade 57, Amsterdam.(0).
Holland.

3. to be the heart, for throughout - Spring Meeting, Canterbury. out of the talle tales (P. mine, duration inclusive (notation and design le The Please read . Please read the meet opwe take the Dr. wedad Tadros and Prof. Dr. J.J. Laarman Dept. of Parasitology, University of Amsterdam
Institute of Tropical Hygiene, dpt. of the Royal Tropical Institute The Netherlands. . TELTHORE NUMBER 020 - 944422 TITLE OF PAPER ople de la informative and succinct as possible. Give generic THE ELUCIDATION OF THE LIFE CYCLE OF SARCOCYSTIS CUNICULI OF THE EUROPEAN RABBIT ORYCTOLAGUS CUNICULUS. RESUME OF PAPER brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be It has recently been demonstrated that the tissue cysts of Sarcocystis parasitizing the musculature of numerous vertebrates, consitute an intermediate stage in the life cycle pattern of a group of endosporulating disporic, tetrazoic eimeriid coccidia. The current communication presents the elucidation of the life cycle of S. cuniculi of the European rabbit, with domestic cats Data will be presented on the light- and electronmicroscopic structure of this sarcocystic species, on the asexual and sexual phases of the life cycle and on serological findings in both the intermediate lagomorph and definitive feline host. (For organiser's use) SESSION TIME DATE PARASE REST. THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED. The elucidation of the life cycle of Sarcocystis cuniculi of the European rabbit Oryctolagus cuniculus.

JAME

Dr. Wedad Tadros & Prof. Dr. J.J. Laarman

ADDRESS

Institute of Tropical Hygiene, Mauritskade 57, Amsterdam.(0).
Holland.

the Street, for three itology - Spring Meeting, Canterbury.

of the other water (19 mins, duration inclusive of disampolon)

read there is a laminy will not be accepted. Please read

. a hole a Prof. Dr. J.J. Laarman and Dr. Wedad Tadros ive ti'l. : i'rof, Dr. gtr.)

I. ACE OF RESPANCE Institute of Tropical Hygiene, dpt.of the Royal Tropical Institute.

Dept. of Parasitology, University of Amsterdam. - TELEPHONE NUMBER 020 - 944422 The Netherlands

4. TUTLE OF PAPER

(plane be an informative and auccinct as possible. Give generic and specific names in full

SOME OBSERVATIONS ON THE EPIDEMIOLOGY, SEROLOGY AND EXPERIMENTAL

TRANSMISSION OF SARCOCYSTIS-INDUCED COCCIDIOSIS IN MAN,

LESUME OF PAPER

(brief but clear resume of contert of paper, not more than 100 words. This will be used to help the organisers but will not be

It is now known that the mature, tetrazoic sporocysts shed by humans and previously referred to as <u>Isospora hominis</u>, represent the end product of the gametogonic development in the human intestinal tract of at least two species of <u>Sarcocystis</u>, namely S.hominis of cattle and S.suihominis of pigs.

Data will be presented on the: (i) natural incidence of S.hominisinduced coccidiosis in selcted samples of human populations from

different geographic, ethnic and age categories; (ii) serological investigations by the I.F.A. for circulating antibodies against Sarcocystis antigen; (iii) experimentally induced cycle of development of S. hominis in a calf and in human volunteers.

(For organiser's use)

SESSI	ON	WEND			
		_TIME	DATI	3	
			 CONTRACTOR OF THE CONTRACTOR	Service distribution of the last of the la	

PLEASE REFERE THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

Some observations on the epidemiology, serology and experimental Transmission of Sarcocystis-induced coccidiosis in man.

JAME

ADDRESS

Prof.Dr. J.J. Laarman & Dr. Wedad Tadros Institute of Tropical Hygiene, Mauritskade 57,

Amsterdam.(0). Holland.

the british Salety for Parabitology - Spring Mee	3.
	ting, Canterbury.
	uration inclusive f discussion)
and a halo on the meet nor	epte . Please read
Orve title: Prof, Dr. etc.)	
Leyden University, Medical	Conton
Parasitology, Rapenburg 33, 071-148333 ext.7465	Leyden, The Netherlands
1. TITLE OF PARED	
(please be as informative and succinct as possi and specific names in full) Clones of axenic Entamoeba histolytica	ble. Give generic
nistorytica nistorytica	
(brief but clear resume of content of paper, not words. This will be used to help the organisers published) By means of a thinly drawn Pasteur pipette single in small volumes of medium in association with fix were compared with the original strain for their toxicity in vitro, and Con A induced agglutination believed that E.histolytica multiplies asexually. Study of clones could give us insight in the gene thamsters will be discussed, as well as the interaction different clones and guinea pig neutrophils.	e amoebae were selected gle amoebae was achieved agellates. These clones growth, virulence, on. It is generally Therefore, the
(For organiser's use)	
SESSION TIME	
AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE	ALLOCATED BELOW
THE Clones of axenic Entered CLEARLY WITHIN THE	BOX PROVIDED.

VAME

Dr.H.J.Bos

Clones of axenic Entamoeba histolytica

ADDRESS

Leyden University, Medical Center, Department of Parasitology, Rapenburg 33, Leyden, the Netherlands

cattle

The British Society for Parasitology - Spring Meeting, Canterbury. 5th - 7th April 1978.

APPLICATION TO PRESENT A FREE PAPER (15 mins. duration inclusive of discussion)

Papers received after 23rd January will not be accepted. Please read enclosed notes on the meeting.

- 1. AUTHOR (S) Dr.F. van Knapen and Dr.S. Fridas
 (Cave title: Prof, Dr. etc.)
- 2. PLACE OF RESEARCH National Institute of Public Health, P.O.Box 1, Bilthoven
 The Netherlands
- 3. TELEPHONE NUMBER (030) 789111
- 4. TITLE OF PAPER
 (please be as informative and succinct as possible. Give generic and specific names in full)

Enzyme-Linked Immunosorbent Assay (ELISA) and its application to the detection

of circulating antigen of Cysticercus bovis in naturally and experimentally infected

5. RESUME OF PAPER
(brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be published)

For the diagnosis of bovine cysticercosis macroscopic examination of carcasses is employed routinely. Serological methods can be of help, provided that they are sensitive, specific, and can preferably be automated. A presentation is given on the usefulness of the Enzyme-Linked Immunosorbent Assay (ELISA) for the serodiagnosis of bovine cysticercosis in the living animal.

Furthermore, the possibility of detecting circulating antigens with the same method (ELISA) in naturally infected animals is shown and the future application of these tests is discussed.

£	(For_organiser's	use	,

SESSION TIME DATE

7. PLEASE REPEAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

TITLE Enzyme-Linked Immunosorbent Assay (ELISA) and its application to the detection of circulating antigen of Cysticercus bovis in naturally and experimentally infected cattle

NAME

F.van Knapen

ADDRESS

National Institute of Public Health, P.O.Box 1, Bilthover. The Netherlands the british Salety for Paramitology - Spring Meeting, Canterbury,

THOR ! THE ENT A FILE PAPER (15 mins. duration inclusive of discussion)

- 1. Al Hoh 1 Dr. J.P. Rotmans cove title: Prof. Dr. etc.)
- 1. ACE OF RESEARCH Leiden University, Medical Center, Department of Parasitology, Rapenburg 33, Leiden, The Netherlands TELEPHONE NUMBER
- 071-148333 ext.7459
- TITLE OF PAPER (please be as informative and succinct as possible. Give generic The use of schistosomal enzymes as antigens in immunodiagnostical tests for schistosomiasis.

RESUME OF PAPER (brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be published)

The purification of a number of schistosomal enzymes (malate dehydrogenase, acidic protease, acetylcholinesterase) has been undertaken, to investigate the use of these enzymes in immunodiagnostical tests (e.g. DASS-test). Purification procedures and biochemical characterization of the enzymes investigated will be discussed. Immunoelectrophoresis experiments and immunization procedures in rabbits with partially purified enzymeantigens, showed that these enzyme preparations are immunologically impure, although considerable purification (compared to the crude homogenate) was obtained. Consequences of these data for the use of these enzymes in immunodiagnostical tests, will be discussed.

,	(For	organi	ser!	s	use)
				-	~ ~ /

SESSION			
 NOTague	TIME		T) A (D) D
35 W A	5 T 15 VA	-	DATE

PLEASE REPEAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

The use of schistosomal enzymes as antigens in immunodiagnostical

VAME

Dr.J.P. Rotmans

ADDRESS

Leiden University, Medical Center, Department of Parasitology, Rapenburg 33, Leiden. The Netherlands.

APPLICATION TO PRESENT A FREE PAPER (15 mins. duration inclusive of discussion)

Papers received after 23rd January will not be accepted. Please read enclosed notes on the meeting.

- 1. AUTHOR (S) Dr. E.J. Ruitenberg, Anneke Elgersma and C.H.J. Lamers (Give title: Prof, Dr. etc.)
- 2. PLACE OF RESEARCH National Institute of Public Health, Bilthoven, The Nether-lands
- 3. TELEPHONE NUMBER 030- 789111 ext. 1075
- 4. TITLE OF PAPER (please be as informative and succinct as possible. Give generic and specific names in full)

Mast cells and eosinophils in Trichinella spiralis infections

5. RESUME OF PAPER

(brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be published)

During the intestinal phase of a T.spiralis infection in both rats and mice in the number of both intestinal mast cells and intra-epithelially located globule leucocytes was observed. The presence of these cells was shown to be T-dependent. Studies involved demonstration of IgE in the intestine

(mast cells and globule leucocytes) and mesenteric lymph nodes (plasma cells).

The origin and the reason for attraction of mast cells to the gut was studied. Next, blood and tissue eosinophils were studied. T.spiralis induced eosinophilia was shown to be thymus-dependent.

The cellular infiltrate will be discussed as immunity-dependent pathological event.

6. (For organiser's use)

SESSION	TIME		DATE
	The state of the s	AND DESCRIPTION OF THE PARTY OF	

7. PLEASE REPEAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

TITLE Mast cells and eosinophils in Trichinella spiralis infections

NAME

Dr. E.J. Ruitenberg

ADDRESS

National Institute of Public Health P.O.Box 1
Bilthoven
The Netherlands

1 1	151	tich Sx	101, 10r	Paramtology	-	Spring	Meeting, Canterbury.
-----	-----	---------	----------	-------------	---	--------	----------------------

ALL CATION TO PRESENT A FILE PAPER	(15 mir	ne. duration of discu	inclusive
------------------------------------	---------	--------------------------	-----------

received after Man danuary will not be accepte. Please read end of d note: on the meet ng.

- Drs.Elisabeth van den Broek (eve title: Prof. Dr. etc.)
- In WE OF RESEARCH University of Amsterdam, Zoological Laboratory, Plantage Doklaan 44, Amsterdam, The Netherlands.
- TELEPHONE NUMBER
- TITLE OF PAPER (please be as informative and succinct as possible. Give generic and specific names in full) Observations on a progenetic stage of Asymphylodora tincae (Trematoda)
- RESUME OF PAPER (brief but clear resume of content of paper, not more than 100 words. This will be used to hel; the organisers but will not be published)

The life cycle of Asymphylodora tincae (Digenea: Monorchiidae) was studied in two small lakes near Amsterdam. The cycle included a progenetic stage, found in Bithynia tentaculata. Field data on the seasonal distribution of this stage will be presented, and the possible influence of the temperature on its appearance will be discussed.

(For organiser's use)

SESSION	OTEMP	
	TIME	DATE

PLEASE REPLA: THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

Observations on a progenetic stage of Asymphylodora tincae (Trematoda)

NAME

ADDRESS

Drs.Elisabeth van den Broek University of Amsterdam, Zoological Laboratory, Plantage Doklaan 44, Amsterdam, The Netherlands.

The fact of the same
th Apr Pros. Throattology - Spring Meeting, Canterbury,
of discussion)
of discussion) with meet ng. 1. a Old of Dr. J.C.M. Vetter T. G.E.
We till: "rof, Dr. ct. JC.M. Klaver-Wesseling & W W.
Dept. of Parasitology, University of Amet.
TELEPHONE NUMBER 020 - 944422 The Netherlands.
(please be as informative and succinct as possible. Give generic ANTIBODIES TO HOOKWORMS.
SECOME OF PAPER (brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be The immune response to hookworm infections has so far received little attention, in contrast to the histopathology. An account will be given of antibody-binding phenomena by all developmental stages, using various methods of treatment of the conjugates.
'. '(For organiser's use)

NOTSENS	TIME	Dage
PLEASE REPEAT THE TETTE OF YOUR		DATE
REASE REPEAT THE TITLE OF YOUR AND WRITE YOUR NAME AND ADDRESS	CLEAR IN THE SPACE	ALLOCATED BELOW
171.1 Antibodies to hookworms	PENETT MITHIN THE	BOX PROVIDED.

IT. Antibodies to hookworms.

VAME

Dr. J.C.M. Vetter, J.C.M. Klaver-Wesseling & W. Visser,

ADDRESS

Institute of Tropical Hygiene, Mauritskade 57, Amsterdam.(0).
Holland.

The British Skiety for Paranitology - Spring Meeting, Canterbury.

AT ACCITION TO PRESENT A FIRE PAPER (15 mins. duration inclusive of discussion)

enclosed on the meet ng.

- 1. A! (!Luk (!) A' Hick (;) Dr.F.H.M.Borgsteede
- IN OF RESEARCH Central Veterinary Institute, Dept. of Parasitology, Edelhertweg 13, Lelystad, the Netherlands
- TELEPHONE NUMBER 03200-21914
- 4. TITLE OF PAPER (please be as informative and succinct as possible. Give generic and specific names in full) Observations on the post-parturient rise in nematode egg-output
- RESUME OF PAPER (brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be

The nematode egg-output of twenty young cows all calving for the first time was followed 3 weeks before calving till 14 weeks afterwards. A marked rise in egg-output in the first week after parturition was observed mainly caused by Ostertagia spp., Trichostrongylus spp. and Cooperia oncophora. The possible reasons for this rise are discussed together with the epidemiological consenquences.

(For organiser's use)

SESSION TIME DATE

PLEASE REPLAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

The Observations on the post-parturient rise in nematode egg-output

VAME

Dr.F.H.M.Borgsteede

ADDRESS

Central Veterinary Institute, Dept.of Parasitology, Edelhertweg 13, Lelystad. the Netherlands

APPLICATION TO PRESENT A FREE PAPER (15 mins. duration inclusive of discussion)

Papers received after 23rd January will not be accepted. Please read enclosed notes on the meeting.

- 1. AUTHOR (S) Drs M. Eysker (Cive title: Prof, Dr. etc.)
- Vet. Parasitology & Parasitic Diseases

 2. PLACE OF RESEARCH State Univ. Utrecht; Yalelaan 7 Utrecht, Uithof, Netherlands
- 3. TELEPHONE NUMBER 030-531114
- 4. TITLE OF PAPER The population dynamics of Trichostrongylus vitrinus and T.colubr(please be as informative and succinct as possible. Give generic and specific names in full)

The population dynamics of Trichostrongylus vitrinus and Trichostrongylus colubriformis in sheep in the Netherlands.

5. RESUME OF PAPER (brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be published)

From 1974 to 1977 the following observations were made on small intestinal Trichostrongylus species of sheep. From November onwards a decline in the number of worms was seen in lambs combined with a swing from the dominant species T. vitrinus to T. colubriformis and occasionally a high percentage of inhibited L3. In barren ewes very high percentages of inhibited L3 were found. In 6 out of 7 ewes harbouring 5th stage male worms, these were T. colubriformis. In lambing ewesdevellopping stages of Trichostrongylus were found and T. vitrinus was more abundant than T. colubriformis.

(, (For organiser's use)

SESSION TIME DATE

7. PLEASE REPEAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

The population dynamics of Trichostrongylus vitrinus and Trichostrongylus colubriformis in sheep in the Netherlands.

VAME

Drs M. Eysker; Institute of veterinary Parasitology and parasitic Diseases; State University of Utrecht; Yalelaan 7, Utrecht, Uithof, Netherlands.

ADDRESS

APPLICATION TO PRESENT A FREE PAPER (15 mins. duration inclusive of discussion)

Tapers received after 23rd January w 11 not be accepted. Please read enclosed notes on the meeting.

- 1. AUTHOR (S)

 (Give title: Prof. Dr. etc.)
- Institute of Veterinary Perasitology and Parasitic
 2. PLACE OF RESEARCHDiseases, State University, Utrecht
- 3. TELEPHONE NUMBER The Netherlands 30-531223
- 4. TITLE OF PAPER

 (please be as informative and succinct as possible. Give generic and specific names in full)

 The significance of the lactation priod number for the

spring rise phenomenon in sheep.

5. RESUME OF PAPER
(brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be published)

The spring rise or post-parturient rise in faecal nematode egg counts is a well known phenomenon in sheep. The literature states that the height of the spring rise decreases with the age of the ewes. The paper gives evidence that the height and the pattern of the spring rise does not depend on age of the ewes but on the number of lactation periods.

The significance of lower egg counts and another pattern, during subsequent lactation periods, for an understanding of the spring rise phenomenon will be discussed.

1	(For organiser	•		
	(Tor organiser	S	use	1
	The state of the s			•

CECCTON	2-00mm to 10-10-10		
SESSION	TIME		DATE
	The state of the s	and the second s	DATE
		Married and Address of the Publisher State of the Party State of the P	

7. PLEASE REPEAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

Spring rise phenomenon in sheep.

VAME

Dr. J. Jansen

ADDRESS

Institute of Veterinary Parasitology
Valelasn 7
University Centre "De Withof" - Thicht - 6. The Mathedands

The British Society for Paramitology - Spring Meet	ing,Canterbury.
--	-----------------

A.I.	at along	PRESENT A FILE PAPER	(15 mine.	duration inclusive
1 11				of dimussion)

- enclosed better on the meeting.
- !. At THUR (:)

 Dr.L.H.M. Blommers and Maggy van Lepnep

 (cive title: Prof, Dr. etc.)
- 2. In ACE OF RESEARCH Leyden University, Medical Center, Department
- of Parasitology, Rapenburg 33, Leiden, The Netherlands
 071-148333 ext. 7453
- 4. TITLE OF PAPER

 (please be as informative and succinct as possible. Give generic and specific names in full)

 A new technique for testing the susceptibility of head lice for insecticides.

(brief but clear resume of content of paper, not more than 100 words. This will be used to help the organisers but will not be

Adults are the life cycle stage recommended currently for the use in insecticide experiments in human lice. Adult head lice (Pediculus humanus capitis de Geer) are hardly suited for these tests which were designed originally for the body louse. The number of available specimens is usually low due to difficulties in collecting and rearing, while control mortalities in the tests are often high from premature death by starvation.

These difficulties may be overcome by the use of nymphs I, immediately after their first blood meal. The young lice are very well suited for various kinds of testing methods, such as residue tests, slide-dipping and topical application. The results of experiments with insecticides of interest to head lice control, will be presented.

organiser's use	t	(For organiser	s	use)
-----------------	---	----------------	---	------

SESSIO	N	TIME		
		TIME	DATE	
ULIE INTION			Prince	Contract of the Contract of th

PLEASE REPEAT THE TITLE OF YOUR PAPER IN THE SPACE ALLOCATED BELOW AND WRITE YOUR NAME AND ADDRESS CLEARLY WITHIN THE BOX PROVIDED.

A new technique for testing the susceptibility of head lice for insecticides.

VAME

Dr.L.H.M. Blommers

ADDRESS

Leyden University,
Medical Center,
Department of Parasitology,
Rapenburg 33,
Leyden. The Netherlands,